

RT² Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

Human Cancer Inflammation & Immunity Crosstalk

Cat. no. 330231 PAHS-181Z

For pathway expression analysis

Format	For use with the following real-time cyclers
RT ² Profiler PCR Array, Format A	Applied Biosystems [®] models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad [®] models iCycler [®] , iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf [®] Mastercycler [®] ep realplex models 2, 2s, 4, 4s; Stratagene [®] models Mx3005P [®] , Mx3000P [®] ; Takara TP-800
RT ² Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT ² Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon [®] , DNA Engine Opticon 2; Stratagene Mx4000 [®]
RT ² Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT ² Profiler PCR Array, Format F	Roche [®] LightCycler [®] 480 (96-well block)
RT ² Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT ² Profiler PCR Array, Format H	Fluidigm [®] BioMark™



Description

The Human Cancer Inflammation & Immunity Crosstalk RT² Profiler PCR Array profiles the expression of 84 key genes involved in mediating communication between tumor cells and the cellular mediators of inflammation and immunity. In addition to epithelial and stromal compartments, the tumor microenvironment contains several cell types of the innate and adaptive immune systems including B and T lymphocytes, dendritic cells, and macrophages. In response to tumor-associated antigens presented via MHC Class I molecules, or to abnormal molecular patterns recognized by Toll-like receptors, the immune system eliminates target cells using a variety of effector enzymes and the engagement of pro-apoptotic signals including TRAIL and FAS ligand. If normal homeostasis is not resolved quickly, a state of chronic inflammation can ensue, including locally increased levels of reactive oxygen and nitrogen species that promote genomic instability. Immune cells produce a variety of cytokines that coordinate the inflammatory response, which is fueled by positive feedback loops commonly involving the STAT and NFκB signaling pathways in tumor cells. The resulting upregulation of antiapoptotic and immunosuppressive factors enables transformed cells to proliferate unchecked by the immune system. During cancer progression, the repertoire of chemokines, cytokines, and growth factors that orchestrates normal immune responses can be commandeered to create an immunosuppressive state that facilitates invasion and metastasis. The genes profiled with this array include mediators and effectors of the cross-talk between tumors and the immune system that influences the course of cancer progression. A set of controls present on each array enables data analysis using the Delta-Delta CT method of relative quantification as well as assessment of reverse transcription performance, genomic DNA contamination, and PCR performance. Using real-time PCR, researchers can easily and reliably analyze the expression of a focused panel of genes involved in cancer inflammation and immune crosstalk with this array.

The RT² Profiler PCR Arrays are intended for molecular biology applications. This product is not intended for the diagnosis, prevention, or treatment of a disease.

For further details, consult the *RT² Profiler PCR Array Handbook*.

Shipping and storage

RT² Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT² Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at -20°C.

Note: Ensure that you have the correct RT² Profiler PCR Array format for your real-time cycler (see table above).

Note: Open the package and store the products appropriately immediately on receipt.



Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT² Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
A	ACKR3	AICDA	BCL2	BCL2L1	CCL18	CCL2	CCL20	CCL21	CCL22	CCL28	CCL4	CCL5
B	CCR1	CCR10	CCR2	CCR4	CCR7	CCR9	CD274	CSF1	CSF2	CSF3	CTLA4	CXCL1
C	CXCL10	CXCL11	CXCL12	CXCL2	CXCL5	CXCL9	CXCR1	CXCR2	CXCR3	CXCR4	CXCR5	EGF
D	EGFR	FASLG	FOXP3	GBP1	GZMA	GZMB	HIF1A	HLA-A	HLA-B	HLA-C	IDO1	IFNG
E	IGF1	IL10	IL12A	IL12B	IL13	IL15	IL17A	IL1A	IL1B	IL2	IL23A	IL4
F	IL6	IL8	IRF1	KITLG	MICA	MICB	MIF	MYC	MYD88	NFKB1	NOS2	PDCD1
G	PTGS2	SPP1	STAT1	STAT3	TGFB1	TLR2	TLR3	TLR4	TNF	TNFSF10	TP53	VEGFA
H	ACTB	B2M	GAPDH	HPRT1	RPLP0	HGDC	RTC	RTC	RTC	PPC	PPC	PPC

Gene table: RT² Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Hs.471751	NM_020311	ACKR3	Chemokine (C-X-C motif) receptor 7
A02	Hs.149342	NM_020661	AICDA	Activation-induced cytidine deaminase
A03	Hs.150749	NM_000633	BCL2	B-cell CLL/lymphoma 2
A04	Hs.516966	NM_138578	BCL2L1	BCL2-like 1
A05	Hs.143961	NM_002988	CCL18	Chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated)
A06	Hs.303649	NM_002982	CCL2	Chemokine (C-C motif) ligand 2
A07	Hs.75498	NM_004591	CCL20	Chemokine (C-C motif) ligand 20
A08	Hs.57907	NM_002989	CCL21	Chemokine (C-C motif) ligand 21
A09	Hs.534347	NM_002990	CCL22	Chemokine (C-C motif) ligand 22
A10	Hs.656904	NM_148672	CCL28	Chemokine (C-C motif) ligand 28
A11	Hs.75703	NM_002984	CCL4	Chemokine (C-C motif) ligand 4
A12	Hs.514821	NM_002985	CCL5	Chemokine (C-C motif) ligand 5
B01	Hs.301921	NM_001295	CCR1	Chemokine (C-C motif) receptor 1
B02	Hs.744542	NM_016602	CCR10	Chemokine (C-C motif) receptor 10
B03	Hs.705362	NM_001123396	CCR2	Chemokine (C-C motif) receptor 2
B04	Hs.184926	NM_005508	CCR4	Chemokine (C-C motif) receptor 4
B05	Hs.370036	NM_001838	CCR7	Chemokine (C-C motif) receptor 7
B06	Hs.225946	NM_006641	CCR9	Chemokine (C-C motif) receptor 9
B07	Hs.521989	NM_014143	CD274	CD274 molecule
B08	Hs.173894	NM_000757	CSF1	Colony stimulating factor 1 (macrophage)
B09	Hs.1349	NM_000758	CSF2	Colony stimulating factor 2 (granulocyte-macrophage)
B10	Hs.2233	NM_000759	CSF3	Colony stimulating factor 3 (granulocyte)
B11	Hs.247824	NM_005214	CTLA4	Cytotoxic T-lymphocyte-associated protein 4
B12	Hs.708652	NM_001511	CXCL1	Chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha)
C01	Hs.632586	NM_001565	CXCL10	Chemokine (C-X-C motif) ligand 10
C02	Hs.632592	NM_005409	CXCL11	Chemokine (C-X-C motif) ligand 11
C03	Hs.522891	NM_000609	CXCL12	Chemokine (C-X-C motif) ligand 12
C04	Hs.75765	NM_002089	CXCL2	Chemokine (C-X-C motif) ligand 2
C05	Hs.89714	NM_002994	CXCL5	Chemokine (C-X-C motif) ligand 5
C06	Hs.77367	NM_002416	CXCL9	Chemokine (C-X-C motif) ligand 9
C07	Hs.194778	NM_000634	CXCR1	Chemokine (C-X-C motif) receptor 1
C08	Hs.846	NM_001557	CXCR2	Chemokine (C-X-C motif) receptor 2
C09	Hs.198252	NM_001504	CXCR3	Chemokine (C-X-C motif) receptor 3
C10	Hs.593413	NM_003467	CXCR4	Chemokine (C-X-C motif) receptor 4
C11	Hs.113916	NM_001716	CXCR5	Chemokine (C-X-C motif) receptor 5
C12	Hs.419815	NM_001963	EGF	Epidermal growth factor
D01	Hs.605083	NM_005228	EGFR	Epidermal growth factor receptor
D02	Hs.2007	NM_000639	FASLG	Fas ligand (TNF superfamily, member 6)
D03	Hs.247700	NM_014009	FOXP3	Forkhead box P3
D04	Hs.62661	NM_002053	GBP1	Guanylate binding protein 1, interferon-inducible
D05	Hs.90708	NM_006144	GZMA	Granzyme A (granzyme 1, cytotoxic T-lymphocyte-associated serine esterase 3)
D06	Hs.1051	NM_004131	GZMB	Granzyme B (granzyme 2, cytotoxic T-lymphocyte-associated serine esterase 1)
D07	Hs.719495	NM_001530	HIF1A	Hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)
D08	Hs.713441	NM_002116	HLA-A	Major histocompatibility complex, class I, A

Position	UniGene	GenBank	Symbol	Description
D09	Hs.77961	NM_005514	HLA-B	Major histocompatibility complex, class I, B
D10	Hs.77961	NM_002117	HLA-C	Major histocompatibility complex, class I, C
D11	Hs.840	NM_002164	IDO1	Indoleamine 2,3-dioxygenase 1
D12	Hs.856	NM_000619	IFNG	Interferon, gamma
E01	Hs.160562	NM_000618	IGF1	Insulin-like growth factor 1 (somatomedin C)
E02	Hs.193717	NM_000572	IL10	Interleukin 10
E03	Hs.673	NM_000882	IL12A	Interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)
E04	Hs.674	NM_002187	IL12B	Interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40)
E05	Hs.845	NM_002188	IL13	Interleukin 13
E06	Hs.602618	NM_000585	IL15	Interleukin 15
E07	Hs.41724	NM_002190	IL17A	Interleukin 17A
E08	Hs.1722	NM_000575	IL1A	Interleukin 1, alpha
E09	Hs.126256	NM_000576	IL1B	Interleukin 1, beta
E10	Hs.89679	NM_000586	IL2	Interleukin 2
E11	Hs.591803	NM_016584	IL23A	Interleukin 23, alpha subunit p19
E12	Hs.73917	NM_000589	IL4	Interleukin 4
F01	Hs.654458	NM_000600	IL6	Interleukin 6 (interferon, beta 2)
F02	Hs.624	NM_000584	IL8	Interleukin 8
F03	Hs.436061	NM_002198	IRF1	Interferon regulatory factor 1
F04	Hs.1048	NM_003994	KITLG	KIT ligand
F05	Hs.130838	NM_000247	MICA	MHC class I polypeptide-related sequence A
F06	Hs.731446	NM_005931	MICB	MHC class I polypeptide-related sequence B
F07	Hs.407995	NM_002415	MIF	Macrophage migration inhibitory factor (glycosylation-inhibiting factor)
F08	Hs.202453	NM_002467	MYC	V-myc myelocytomatosis viral oncogene homolog (avian)
F09	Hs.82116	NM_002468	MYD88	Myeloid differentiation primary response gene (88)
F10	Hs.618430	NM_003998	NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1
F11	Hs.709191	NM_000625	NOS2	Nitric oxide synthase 2, inducible
F12	Hs.158297	NM_005018	PDCD1	Programmed cell death 1
G01	Hs.196384	NM_000963	PTGS2	Prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase)
G02	Hs.313	NM_000582	SPP1	Secreted phosphoprotein 1
G03	Hs.743244	NM_007315	STAT1	Signal transducer and activator of transcription 1, 91kDa
G04	Hs.463059	NM_003150	STAT3	Signal transducer and activator of transcription 3 (acute-phase response factor)
G05	Hs.645227	NM_000660	TGFB1	Transforming growth factor, beta 1
G06	Hs.519033	NM_003264	TLR2	Toll-like receptor 2
G07	Hs.657724	NM_003265	TLR3	Toll-like receptor 3
G08	Hs.174312	NM_138554	TLR4	Toll-like receptor 4
G09	Hs.241570	NM_000594	TNF	Tumor necrosis factor
G10	Hs.478275	NM_003810	TNFSF10	Tumor necrosis factor (ligand) superfamily, member 10
G11	Hs.740601	NM_000546	TP53	Tumor protein p53
G12	Hs.73793	NM_003376	VEGFA	Vascular endothelial growth factor A
H01	Hs.520640	NM_001101	ACTB	Actin, beta
H02	Hs.534255	NM_004048	B2M	Beta-2-microglobulin
H03	Hs.544577	NM_002046	GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
H04	Hs.412707	NM_000194	HPRT1	Hypoxanthine phosphoribosyltransferase 1
H05	Hs.546285	NM_001002	RPLP0	Ribosomal protein, large, P0
H06	N/A	SA_00105	HGDC	Human Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

Related products

For optimal performance, RT² Profiler PCR Arrays should be used together with the RT² First Strand Kit for cDNA synthesis and RT² SYBR[®] Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT ² First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT ² SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT ² SYBR Green ROX™ qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT ² SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

* Larger kit sizes available; please inquire.

RT² Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at www.qiagen.com or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

www.qiagen.com

Canada ■ 800-572-9613

Ireland ■ 1800 555 049

Norway ■ 800-18859

China ■ 8621-3865-3865

Italy ■ 800-787980

Singapore ■ 1800-742-4368

Denmark ■ 80-885945

Japan ■ 03-6890-7300

Spain ■ 91-630-7050

Australia ■ 1-800-243-800

Finland ■ 0800-914416

Korea (South) ■ 080-000-7145

Sweden ■ 020-790282

Austria ■ 0800/281010

France ■ 01-60-920-930

Luxembourg ■ 8002 2076

Switzerland ■ 055-254-22-11

Belgium ■ 0800-79612

Germany ■ 02103-29-12000

Mexico ■ 01-800-7742-436

UK ■ 01293-422-911

Brazil ■ 0800-557779

Hong Kong ■ 800 933 965

The Netherlands ■ 0800 0229592

USA ■ 800-426-8157



Sample & Assay Technologies